

★ May ★

★ is ★

BHSM!!

Fluency

★ Facts ★

#1

A fluency disorder is a speech disorder characterized by deviations in continuity, smoothness, rhythm, and/or effort with which phonologic, lexical, morphologic, and/or syntactic language units are spoken

#2


When recovery occurs, it is likely to do so from 6 to 36 months after onset, with most children recovering within the first 1 to 2 years after their stuttering was first noted

#3

Incidence of childhood stuttering is highest between a child's second and fourth birthdays, ultimately affecting 4% to 5% of the population

#4

There is no known cause or cure for stuttering although more research is being done to help us better understand the neurological and environmental factors that may contribute to this speech disorder.



★ statistics from www.asha.org

★ May ★

★ is ★

BHSM!!

Fluency

★ Facts ★

#1

A fluency disorder is a speech disorder characterized by deviations in continuity, smoothness, rhythm, and/or effort with which phonologic, lexical, morphologic, and/or syntactic language units are spoken

#2


When recovery occurs, it is likely to do so from 6 to 36 months after onset, with most children recovering within the first 1 to 2 years after their stuttering was first noted

#3

Incidence of childhood stuttering is highest between a child's second and fourth birthdays, ultimately affecting 4% to 5% of the population

#4

There is no known cause or cure for stuttering although more research is being done to help us better understand the neurological and environmental factors that may contribute to this speech disorder.



★ statistics from www.asha.org

★ May ★

★ is ★

BHSM!!

Fluency

★ Facts ★

#1

A fluency disorder is a speech disorder characterized by deviations in continuity, smoothness, rhythm, and/or effort with which phonologic, lexical, morphologic, and/or syntactic language units are spoken

#2


When recovery occurs, it is likely to do so from 6 to 36 months after onset, with most children recovering within the first 1 to 2 years after their stuttering was first noted

#3

Incidence of childhood stuttering is highest between a child's second and fourth birthdays, ultimately affecting 4% to 5% of the population

#4

There is no known cause or cure for stuttering although more research is being done to help us better understand the neurological and environmental factors that may contribute to this speech disorder.



★ statistics from www.asha.org